



Alere Afinion™ HbA1c

Test for quantitative determination of glycated haemoglobin (HbA1c) in human whole blood, used to monitor metabolic control in patients with diabetes. May be used for diagnosis of diabetes and in identifying patients who may be at risk for developing diabetes, when in compliance with national regulations.*

Simple. Fast. Reliable.

- 3 minutes test time
- Excellent precision
- No interference from Hb variants
- CLIA waived

*May only be used for monitoring in USA.



TAP HERE TO SEE THE PRODUCT



The 3-minute Alere Afinion HbA1c testing fits perfectly within the workflow

“Immediate feedback motivates patients to make the necessary treatment changes.”

Cagliero E et. al. Diabetes Care 1999;22:1785-1789.

“Use of point-of-care testing for HbA1c allows for timely decisions on therapy changes, when needed.”

ADA, 2011. Diabetes Care 2011;34 (supl 1):s11-s61.

3-step procedure

1



2



3



Facts

- 1.5 µL sample volume.
- Configurable display of HbA1c units.
- 4-15% HbA1c, 20-140 mmol/mol HbA1c.
- 3 months room temperature storage.
- Ready to use Controls in 2 levels.
- NGSP and IFCC certified.

Other available tests

Alere Afinion ACR, Alere Afinion CRP and Alere Afinion Lipid Panel



Alere Technologies AS
Kjelsåsveien 161
P.O. Box 6863 Rodeløkka
NO-0504 Oslo, Norway
alere.com

